

WHAT CAN YOU DO?

- Speak slowly and clearly
- Use short sentences
- Keep calm and friendly
- Remind them of the date, time and place
- Bring personal belongings such as glasses, hearing aids, dentures, toiletries, perfume, shaving gel
- Create familiar surroundings
- Avoid too many visitors
- Do not be too demanding or impatient
- Avoid conflict and long discussions
- Take part in a diary



TALK TO US, SHARE YOUR OBSERVATIONS WITH US. THE ACUTE CONFUSIONAL STATE IS REVERSIBLE AND TRANSIENT. REPERCUSSIONS ARE RARE.

Hôpitaux Robert Schuman

9, Rue Edward Steichen, L-2540 Luxembourg
T +352 286 911
www.facebook.com/hopitauxrobertschuman
www.hopitauxschuman.lu

HOPITAL KIRCHBERG'S INTENSIVE CARE UNIT

9, rue Edward Steichen
L-2540 Luxembourg
Tél : (+352) 286 2 5370
Fax : (+352) 286 2 5347

VISITING HOURS :

de 12h30 à 13h30
de 18h00 à 19h30

ZITHAKLINIK'S INTENSIVE CARE UNIT

36, rue de Ste Zithe
L-2763 Luxembourg
Tél : (+352) 286 3 4540
Fax : (+352) 286 3 5920

VISITING HOURS :

de 12h30 à 13h00
de 18h00 à 19h00

DELIRIUM



**ACUTE
CONFUSIONAL
STATE**

WHAT IS AN ACUTE CONFUSIONAL STATE ?

An acute confusional state is a sudden onset of consciousness and concentration troubles.

WARNING SIGNS AND SYMPTOMS:

- Disorientation in time and space
- Lack of concentration
- Delayed reactions
- Hallucinations
- Anxiety
- Aggressiveness
- Apathy
- Agitation
- Lack of sleep/insomnia
- Abnormal and inappropriate stereotypical movements



PROBABLE CAUSES :

One or more triggers such as:

- Infection
- Pain
- Withdrawal (alcohol, cigarettes, medication...)
- Medication
- Unusual surroundings
- Surgery
- State after resuscitation
- Excess of Stimuli (light, noise, catheters, monitoring...)
- Immobility
- Difficulty in communication (language, hearing aids, glasses...)
- Dehydration
- Stress

WHO IS AT RISK OF DEVELOPING AN ACUTE CONFUSIONAL STATE?

In principle: everyone

Favourable factors : old age and dementia

WHAT CAN WE DO?

- Prevention
- Treat the causes
- Early recognition
- Elimination of disturbances (Day/Night rhythm, noise...)
- Encourage Orientation in time and place, propose clues (clock, calendar...)
- Implement medication if necessary
- Pain treatment
- Encourage fast mobility recovery



WHAT TO AVOID AS MUCH AS POSSIBLE?

- Selfharm and foreign injuries
- Additional medication
- Immobilization of limbs and body